

Characterization of edge-colored complete graphs with properly colored Hamilton paths

Jinfeng Feng* Hans-Erik Giesen† Yubao Guo‡ Gregory Gutin§
Tommy Jensen¶ Arash Rafiey||

Abstract

An edge-colored graph H is properly colored if no two adjacent edges of H have the same color. In 1997, J. Bang-Jensen and G. Gutin conjectured that an edge-colored complete graph G has a properly colored Hamilton path if and only if G has a spanning subgraph consisting of a properly colored path C_0 and a (possibly empty) collection of properly colored cycles C_1, C_2, \dots, C_d such that $V(C_i) \cap V(C_j) = \emptyset$ provided $0 \leq i < j \leq d$. We prove this conjecture.

Keywords: edge-colored graphs, properly colored Hamilton paths.

1 Introduction

Let $G = (V, E)$ be a complete graph, and let $c : E \rightarrow \{1, 2, \dots, \chi\}$ be a fixed (not necessarily proper) edge-coloring of G with χ colors, $\chi \geq 2$. With given c , G is called a χ -edge-colored (or, *edge-colored*) complete graph. A subgraph $H \subseteq G$ is called *properly colored* if c defines a proper edge-coloring of H .

The existence of properly colored Hamilton paths and cycles has been studied in several papers; this topic was surveyed in [2] and later in Chapter 11 of [3]. While there are characterizations [6, 11] (see also Chapter 11 of [3]) of 2-edge-colored complete graphs with properly colored Hamilton cycles, no such characterization is known for χ -edge-colored complete graphs with $\chi \geq 3$, and it is still an open question to determine the computational complexity of this problem [8].

*Lehrstuhl C fuer Mathematik, RWTH Aachen University, 52056 Aachen, Germany.

†Lehrstuhl C fuer Mathematik, RWTH Aachen University, 52056 Aachen, Germany.

‡Lehrstuhl C fuer Mathematik, RWTH Aachen University, 52056 Aachen, Germany.

§Corresponding author. Department of Computer Science, Royal Holloway, University of London, Egham, Surrey, TW20 0EX, UK. Gutin@cs.rhul.ac.uk

¶Institut fuer Mathematik, Universitat Klagenfurt, Universitätsstrasse 65-67, 9020 Klagenfurt, Austria.

||Department of Computer Science, Royal Holloway, University of London, Egham, Surrey, TW20 0EX, UK.

The most studied *possibly* sufficient condition for an edge-colored complete graph with n vertices to have a properly colored Hamilton cycle is $\Delta_{mon} < \lfloor n/2 \rfloor$, where Δ_{mon} is the maximal number of edges of the same color incident to the same vertex. This was conjectured by B. Bollobás and P. Erdős [9] in 1976, but remains unsolved. The best result so far for 'small' values of n is by J. Shearer [12]: $7\Delta_{mon} < \lfloor n/2 \rfloor$ guarantees the existence of a properly colored Hamilton cycle. The best result so far for large values of n is due to N. Alon and G. Gutin [1]: For every $\epsilon > 0$ and $n = n_\epsilon$ large enough, $\Delta_{mon} \leq (1 - \frac{1}{\sqrt{2}} - \epsilon)\lfloor n/2 \rfloor$ implies the existence of a properly colored Hamilton cycle.

For the case of properly colored Hamilton paths, the situation is somewhat different. Let the abbreviation PCHP stand for "properly colored Hamilton path". Let G be an edge-colored graph. A *properly colored cycle factor* of G is a spanning subgraph of G consisting of properly colored cycles C_1, C_2, \dots, C_d such that $V(C_i) \cap V(C_j) = \emptyset$ provided $1 \leq i < j \leq d$. A *properly colored 1-path-cycle factor* of G is a spanning subgraph of G consisting of a properly colored path C_0 and a (possibly empty) collection of properly colored cycles C_1, C_2, \dots, C_d such that $V(C_i) \cap V(C_j) = \emptyset$ provided $0 \leq i < j \leq d$.

The following theorem gives a PCHP characterization for the case of just two colors:

Theorem 1.1 [2] *A 2-edge-colored complete graph G has a PCHP if and only if G contains a properly colored 1-path-cycle factor.*

It is conjectured in [2] that the above theorem holds for any number of colors. We call it the BJG conjecture. In support of the BJG conjecture, the following result was proved in [5]: If a χ -edge-colored complete graph G ($\chi \geq 2$) contains a properly colored cycle factor, then G contains a PCHP.

It is easy to see that the BJG conjecture in [2] can be reduced to the following:

Conjecture 1.2 (PCHP Conjecture) *Let $\chi \geq 3$ and let G be a χ -edge-colored complete graph. Assume that there exist $C, P \subseteq G$, where C is a properly colored cycle and P a properly colored path, such that $V(C) \cap V(P) = \emptyset$ and $V(C) \cup V(P) = V(G)$. Then G contains a PCHP.*

In this paper we prove the PCHP conjecture and, thus, the BJG conjecture. Since it takes polynomial time to check whether an edge-colored graph has a properly colored 1-path-cycle factor [2], our result implies that the PCHP problem is polynomial time solvable for edge-colored complete graphs. The proof of Theorem 2.1 is constructive and can be turned into a polynomial time algorithm for transforming a properly colored 1-path-cycle factor into a properly colored Hamilton path.

This gives, in particular, some indication that the problem of the existence of a properly colored Hamilton cycle in an edge-colored graph may be polynomial time solvable after all.

The situation may remind one of that with the existence of Hamilton paths and cycles in semicomplete multipartite digraphs (SMDs) [4] (see also Chapter 5 in [3]). Both Hamilton path and cycle problems for SMDs are polynomial time solvable, but only for the Hamilton path problem we have a nice characterization (see, e.g., [10] or Chapter 5 in [3]) so far.

In passing we mention a simple sufficient condition proved in [7] for the existence of a PCHP in an edge-colored K_n : K_n has no monochromatic triangles.

2 Results

If H is connected, the *distance in H* between two vertices $u, v \in V(H)$ is the length of a shortest path in H from u to v , and we denote it by $\text{dist}_H(u, v)$.

Theorem 2.1 *The PCHP Conjecture holds.*

Proof: Let $C = v_1 \dots v_n v_1$ ($n \geq 3$) and $P = u_1 \dots u_m$ ($m \geq 1$). Throughout we will perform addition and subtraction in the indices of the vertices $v_j \in C$ modulo n .

Let $j \in \{1, 2, \dots, n\}$. If $m \geq 2$ and $c(u_1 v_j) \neq c(u_1 u_2)$, then at least one of the paths $u_m u_{m-1} \dots u_1 v_j v_{j+1} \dots v_{j-1}$ and $u_m u_{m-1} \dots u_1 v_j v_{j-1} \dots v_{j+1}$ is a PCHP. Similarly there exists a PCHP if $c(u_m v_j) \neq c(u_{m-1} u_{m-2})$. So we may assume the following:

(1) If $m \geq 2$, then $c(u_1 v_j) = c(u_1 u_2)$ and $c(u_m v_j) = c(u_{m-1} u_m)$ for every $j = 1, 2, \dots, n$.

Thus, to complete the proof of this theorem it suffices to prove the following claim:

Claim A *If (1) is satisfied, then there exists a PCHP H in G with u_1 as its first vertex, such that the initial edge of H is either $u_1 u_2$ or one of the edges $u_1 v_j$ ($1 \leq j \leq n$), and such that if $m \geq 2$, then u_m is the last vertex of H and the last edge of H is either $u_{m-1} u_m$ or one of the edges $v_j u_m$ ($1 \leq j \leq n$).*

Let $b(P, C) = 2(n - 3) + m$; we notice that $b(P, C) \geq 1$. Suppose that Claim A is false, and let (G, P, C, c) be a counterexample with a minimal value of $b(P, C)$.

If $m = 1$, then either $u_1 v_1 v_2 \dots v_n$ or $u_1 v_1 v_n \dots v_2$ is a PCHP as desired. Thus, we have established $m \geq 2$ and $b(P, C) \geq 2$.

Now we prove

(2) $m \geq 3$.

Suppose that $m = 2$. With $x = c(u_1u_2)$, we have $c(u_1v_j) = c(u_2v_j) = x$ for all $j = 1, 2, \dots, n$, by (1). Choose r such that $c(v_{r-1}v_r) \neq x \neq c(v_{r+1}v_{r+2})$; this is possible if r can be chosen with $c(v_rv_{r+1}) = x$, and otherwise it is trivial. But then the path $u_1v_rv_{r-1} \dots v_{r+2}v_{r+1}u_2$ yields a contradiction.

We continue to prove further properties of the coloring c .

(3) For every i , $1 < i < m$, there exist r and s ($1 \leq r, s \leq n$), such that $c(u_iu_{i+1}) \neq c(u_iv_r)$, $c(u_{i-1}u_i) \neq c(u_iv_s)$.

Otherwise the path $P' = P - u_1 - u_2 - \dots - u_{i-1}$ satisfies (1) in $G' = G - u_1 - u_2 - \dots - u_{i-1}$ (with u_i in place of u_1). Since $b(P', C) < b(P, C)$, there exists a PCHP H' in G' that starts from u_i with one of the edges u_iu_{i+1} or u_iv_j , $1 \leq j \leq n$ and finishes in u_m with $u_{m-1}u_m$ or with an edge v_kv_m , $1 \leq k \leq n$. Then $u_1 \dots u_{i-1}H'$ is a PCHP in G of the desired type having u_1u_2 as its initial edge, a contradiction. A similar argument shows the existence of s .

Suppose that $c(v_{i-1}v_i) = c(v_jv_{j+1}) = x$, where $\text{dist}_C(v_i, v_{j+1}) \geq 3$. If $c(v_iv_j) = x$, then define $G' = G - v_i - v_{i+1} - \dots - v_j$ and define $c' : E(G') \rightarrow \{1, 2, \dots, \chi\}$ by

$$c'(e) = \begin{cases} x & \text{if } e = v_{i-1}v_{j+1} \\ c(e) & \text{otherwise.} \end{cases}$$

Both P and the cycle $C' = v_{j+1}v_{j+2} \dots v_nv_1 \dots v_{i-1}v_{j+1}$ are properly colored by c' . Moreover, P clearly satisfies (1) with respect to C' . Since $b(P, C') < b(P, C)$, there exists a PCHP P_1 in G' with initial edge u_1u_2 or u_1v_r for some $r \in \{j+1, j+2, \dots, n, 1, \dots, i-1\}$. If $v_{i-1}v_{j+1} \in P_1$, then we find the desired PCHP H in G by replacing the edge $v_{i-1}v_{j+1}$ by the path $v_{i-1}v_i \dots v_jv_{j+1}$. Otherwise, if $v_{i-1}v_{j+1} \notin P_1$, then P_1 is properly colored in G and satisfies (1) with respect to $C'' = v_iv_{i+1} \dots v_jv_i$, which is a properly colored cycle. By $b(P_1, C'') < b(P, C)$, the desired PCHP exists in G , a contradiction.

Thus, we have the following:

(4) Assume that $\text{dist}_C(v_s, v_{t+1}) \geq 3$. If $c(v_{s-1}v_s) = c(v_tv_{t+1}) = x$, then $c(v_sv_t) \neq x$.

Consider the path $u_1 \dots u_{p-2}u_{p-1}v_{q-1}v_{q-2} \dots v_{q+1}v_qu_pu_{p+1} \dots u_m$. As it cannot be a PCHP we conclude the following.

(5) Let $2 \leq p \leq m$ and $1 \leq q \leq n$. Then at least one of the following holds.




Figure 1: The path used to prove (5).

- a) $p \geq 3$ and $c(u_{p-2}u_{p-1}) = c(u_{p-1}v_{q-1})$
- b) $c(u_{p-1}v_{q-1}) = c(v_{q-2}v_{q-1})$
- c) $c(u_p v_q) = c(v_q v_{q+1})$
- d) $p < m$ and $c(u_p v_q) = c(u_p u_{p+1})$.

Considering the path $u_1 \dots u_{p-2} u_{p-1} v_{q+1} v_{q+2} \dots v_{q-1} v_q u_p u_{p+1} \dots u_m$ similarly leads to:

(6) Let $2 \leq p \leq m$ and $1 \leq q \leq n$. Then at least one of the following holds.

- a) $p \geq 3$ and $c(u_{p-2}u_{p-1}) = c(u_{p-1}v_{q+1})$
- b) $c(u_{p-1}v_{q+1}) = c(v_{q+1}v_{q+2})$
- c) $c(u_p v_q) = c(v_{q-1}v_q)$
- d) $p < m$ and $c(u_p v_q) = c(u_p u_{p+1})$.

In several of the following applications of (5) and (6) it will be useful to note that (5c) and (6c) are mutually exclusive statements for any values of p and q , since C is properly colored, and that (5d) and (6d) are identical statements.

For the remaining part of the paper we define $x = c(u_1 u_2)$ and $y = c(u_{m-1} u_m)$.

(7) Assume $c(u_2 v_j) = z \neq c(u_2 u_3)$ for some $j \in \{1, 2, \dots, n\}$.




Figure 2: The path used to prove (6).

- a) If $c(v_j v_{j+1}) \neq z$ then $c(v_{j-2} v_{j-1}) = x$, and if $c(v_{j-1} v_j) \neq z$ then $c(v_{j+1} v_{j+2}) = x$.**
- b) If $z = x$ or $c(v_{j-1} v_j) \neq z \neq c(v_j v_{j+1})$, then $c(v_{j-2} v_{j-1}) = c(v_{j+1} v_{j+2}) = c(v_{j-1} v_{j+1}) = x$ and $n \in \{3, 5\}$.**

If $c(v_{j-1} v_j) \neq z$, then $c(v_{j+1} v_{j+2}) = x$ follows from (6) with $p = 2$ and $q = j$ (only (b) of (6) is not necessarily false). If $c(v_j v_{j+1}) \neq z$, then $c(v_{j-2} v_{j-1}) = x$ similarly follows from (5). This shows (a).

Now assume $c(v_{j-1} v_j) \neq z \neq c(v_j v_{j+1})$ or $z = x$. In the case $z = x$ the fact that c is a proper coloring of C together with (a) implies $c(v_{j-2} v_{j-1}) = c(v_{j+1} v_{j+2}) = x$. The same conclusion follows directly from (a) when $c(v_{j-1} v_j) \neq z \neq c(v_j v_{j+1})$. By symmetry, and since $c(v_{j-1} v_j) \neq c(v_j v_{j+1})$, we may assume $c(v_{j-1} v_{j+1}) \neq c(v_j v_{j+1})$. Since the Hamilton path $u_1 v_{j+2} v_{j+3} \dots v_{j-2} v_{j-1} v_{j+1} v_j u_2 \dots u_m$ fails to be a PCHP, it follows that $c(v_{j-1} v_{j+1}) = x$.

Consider (4) for $s = j - 1$ and $t = j + 1$. The conclusion $c(v_s v_t) \neq x$ of (4) implies $\text{dist}_C(v_{j-1}, v_{j+2}) \leq 2$, which is only possible for $n \leq 5$. Moreover, the edges $v_{j-2} v_{j-1}$ and $v_{j+1} v_{j+2}$, both of color x , are not adjacent on C , which implies $n \neq 4$. Thus (b) is proved.

(8) Assume $c(u_{m-1} v_k) = z \neq c(u_{m-2} u_{m-1})$ for some $k \in \{1, 2, \dots, n\}$.

- a) If $c(v_{k-1} v_k) \neq z$ then $c(v_{k+1} v_{k+2}) = y$, and if $c(v_k v_{k+1}) \neq z$ then $c(v_{k-2} v_{k-1}) = y$.**
- b) If $z = y$ or $c(v_{k-1} v_k) \neq z \neq c(v_k v_{k+1})$, then $c(v_{k-2} v_{k-1}) = c(v_{k+1} v_{k+2}) = c(v_{k-1} v_{k+1}) = y$ and $n \in \{3, 5\}$.**

The proof of (8) is similar to that of (7).

(9) Assume $c(u_2v_j) \neq x$ for all $j = 1, 2, \dots, n$. Then there exists $j \in \{1, 2, \dots, n\}$ such that

a) $c(u_2v_{j-1}) = c(u_2v_j) \neq c(u_2u_3)$, and

b) $c(v_{j-2}v_{j-1}) = c(v_jv_{j+1}) = x$.

By (3) there exists $j \in \{1, 2, \dots, n\}$ such that $c(u_2v_j) = z \neq c(u_2u_3)$. We may assume $c(v_jv_{j+1}) \neq z$ (if not, then $c(v_{j-1}v_j) \neq z$ holds, and we may renumber the vertices on C so that $v_{j+\ell}$ becomes $v_{j-\ell}$ for all $\ell = 0, 1, 2, \dots$ without change of the conclusion). Then (7a) implies $c(v_{j-2}v_{j-1}) = x$.

Again by (3) there exists $r \in \{1, 2, \dots, n\}$ such that if $m \geq 4$, then $c(u_3v_r) \neq c(u_3u_4)$ holds.

Suppose $c(u_2v_{j-1}) \neq z$. Since $u_1v_{r+1}v_{r+2} \dots v_{j-2}v_{j-1}u_2v_jv_{j+1} \dots v_{r-1}v_ru_3u_4 \dots u_m$ is not a PCHP, at least one of the following holds:

i) $c(u_1v_{r+1}) = c(v_{r+1}v_{r+2})$,

ii) $c(u_3v_r) = c(v_{r-1}v_r)$,

iii) $r = j$ and $c(u_2v_r) = c(u_3v_r)$.

Since $u_1v_{r-1}v_{r-2} \dots v_{j+1}v_ju_2v_{j-1}v_{j-2} \dots v_{r+1}v_ru_3u_4 \dots u_m$ is not a PCHP, at least one of the following holds:

iv) $c(u_1v_{r-1}) = c(v_{r-2}v_{r-1})$,

v) $c(u_3v_r) = c(v_rv_{r+1})$,

vi) $r = j - 1$ and $c(u_2v_r) = c(u_3v_r)$.

Let $p = 3$ and $q = r$, and observe that neither of (5a), (5d), (6a) or (6d) holds.

We will now show that (iii) and (vi) do not hold. If $r = j$, then $c(v_{r-2}v_{r-1}) = x \neq c(u_2v_{r-1})$, hence also (5b) does not hold, and (5c) must be satisfied, that is, $c(u_3v_r) = c(v_rv_{r+1})$. In particular, if $r = j$, then $c(u_3v_r) = c(v_jv_{j+1}) \neq z = c(u_2v_j)$, contrary to (iii).

If $r = j - 1$, then $c(u_2v_{r+1}) = c(u_2v_j) = z$ and $c(v_{r+1}v_{r+2}) = c(v_jv_{j+1}) \neq z$, so (6b) does not hold. Then (6c) implies $c(u_3v_r) = c(v_{r-1}v_r) = c(v_{j-2}v_{j-1}) = x$. In particular, if $r = j - 1$, then $c(u_2v_r) \neq c(u_3v_r)$ follows, hence also (vi) does not hold.

We deduce that (i) or (ii) is true, and that (iv) or (v) is true. Now (6c) is equivalent to (ii), and (6b) and (i) are not both true (by (1) and our assumption), therefore (ii) holds. Similarly (5c) is equivalent to (v), and (5b) contradicts (iv), so also (v) holds. But (ii) contradicts (v), since C is properly colored. This establishes $c(u_2v_{j-1}) = z$.

Finally $c(v_jv_{j+1}) = x$ follows from (7a).

(10) Assume $c(u_{m-1}v_j) \neq y$ for all $j = 1, 2, \dots, n$. Then there exists $k \in \{1, 2, \dots, n\}$ such that

- a) $c(u_{m-1}v_{k-1}) = c(u_{m-1}v_k) \neq c(u_{m-2}u_{m-1})$, and
- b) $c(v_{k-2}v_{k-1}) = c(v_kv_{k+1}) = y$.

The proof is similar to the proof of (9).

(11) Assume $z = c(u_2v_j) = c(u_2v_{j-1}) \notin \{x, c(u_2u_3)\}$ for some $j \in \{1, 2, \dots, n\}$. Furthermore assume $c(v_jv_{j+1}) \neq z$. Then one of the following holds:

- a) n is an even number, and $n/2$ edges of C have color x .
- b) If $c(v_{j'}v_{j'+1}) = x$, then $v_{j'} \in S = \{v_{j-4}, v_{j-2}, v_j, v_{j+2}\}$.

First $c(v_{j-2}v_{j-1}) = c(v_jv_{j+1}) = x$ follows from two applications of (7a), applying (7) to u_2v_j and u_2v_{j-1} in this order. Since C is properly colored, we conclude that $c(v_{j'}v_{j'+1}) = x$ implies $v_{j'} \notin \{v_{j-3}, v_{j-1}, v_{j+1}\}$. Assume that (b) does not hold, and choose any $v_{j'} \in V(C) \setminus S$ satisfying $c(v_{j'}v_{j'+1}) = x$. Then $n \geq 8$ follows from $v_{j'} \notin S \cup \{v_{j-3}, v_{j-1}, v_{j+1}\}$. We will prove the following statement.

$$(*) \quad c(v_{j'-2}v_{j'-1}) = c(v_{j'+2}v_{j'+3}) = x.$$

First we suppose $c(v_{j'-2}v_{j'-1}) \neq x$. Then $v_{j'} \notin S \cup \{v_{j-3}, v_{j-1}, v_{j+1}\}$ and $n \geq 8$ imply $\text{dist}_C(v_{j-1}, v_{j'+1}) \geq 3$. By (4) with $s = j - 1$ and $t = j'$ we have $c(v_{j-1}v_{j'}) \neq x$. We consider the path $P_1 = u_1v_{j'-1}v_{j'-2} \dots v_{j+1}v_ju_2u_3 \dots u_m$ and the cycle $C_1 = v_{j'}v_{j'+1} \dots v_{j-2}v_{j-1}v_{j'}$, which are properly colored and satisfy (1). Since $b(C_1, P_1) < b(C, P)$ holds, our minimality assumption yields a PCHP as in Claim A, which is a contradiction. So $c(v_{j'-2}v_{j'-1}) = x$ holds. Now suppose $c(v_{j'+2}v_{j'+3}) \neq x$. Then by (4) with $s = j$ and $t = j' + 1$ we similarly have $c(v_jv_{j'+1}) \neq x$, and we consider the path $P_2 = u_1v_{j'+2}v_{j'+3} \dots v_{j-2}v_{j-1}u_2u_3 \dots u_m$ and the cycle $C_2 = v_{j'+1}v_{j'} \dots v_{j+1}v_jv_{j'+1}$ instead, again with a contradiction. Thus we have also $c(v_{j'+2}v_{j'+3}) = x$, which finishes the proof of (*).

Applying (*) recursively, it follows that $c(v_{j'+2\ell}v_{j'+2\ell+1}) = x$ holds for every $\ell \in \mathbb{N}$ with $v_{j'+2\ell} \notin S \cup \{v_{j-3}, v_{j-1}, v_{j+1}\}$. In particular, either $c(v_{j+3}v_{j+4}) = x$ or $c(v_{j+4}v_{j+5}) = x$ holds ($v_{j+3} \notin S$ follows from $n \geq 8$, and $v_{j+4} \in S$ only occurs if $n = 8$ and $v_{j'} = v_{j+3}$, in which case $c(v_{j+3}v_{j+4}) = x$ follows). However, applying (*), $c(v_{j+3}v_{j+4}) = x$ would imply $c(v_{j+1}v_{j+2}) = x$, contradicting the fact that C is properly colored. So $c(v_{j+4}v_{j+5}) = x$ holds. Similar reasoning leads to $c(v_{j-6}v_{j-5}) = x$. It follows from (*) that all of $v_{j+4}v_{j+5}, v_{j+6}v_{j+7}, \dots, v_{j-6}v_{j-5}$ are colored x .

Since both $v_{j+4}v_{j+5}$ and $v_{j-6}v_{j-5}$ are colored x , it follows from (*) that $v_{j+2}v_{j+3}$ and $v_{j-4}v_{j-3}$ are also colored x . Combining this with the fact that v_jv_{j+1} and $v_{j-2}v_{j-1}$ are both colored x , we have shown that (a) is true, which proves (11).

(12) There is an index j , $1 \leq j \leq n$, such that $c(u_2v_j) = x$ or $c(u_{m-1}v_j) = y$.

Suppose $c(u_2v_j) \neq x$ and $c(u_{m-1}v_j) \neq y$ for all $j = 1, 2, \dots, n$. By (9) and (10) we may choose $j, k \in \{1, 2, \dots, n\}$ such that j satisfies (9a) and (9b), and k satisfies (10a) and (10b).

By (9b) we have $c(v_{j-2}v_{j-1}) = c(v_jv_{j+1}) = x$, from which $m \geq 4$ follows, as otherwise $u_1u_2v_jv_{j+1} \dots v_{j-2}v_{j-1}u_3$ would be a PCHP. Now the path

$$u_1v_{k-1}v_{k-2} \dots v_{j+1}v_ju_2u_3 \dots u_{m-2}u_{m-1}v_kv_{k+1} \dots v_{j-2}v_{j-1}u_m$$

is not a PCHP, so $y = x$ follows.

We will show that (11a) holds. So suppose not; then it follows from (11) that (11b) holds. By (10b) we have $c(v_{k-2}v_{k-1}) = c(v_kv_{k+1}) = x$, which by (11b) implies $v_k \in \{v_{j-2}, v_j, v_{j+2}\}$.

The case $v_k = v_j$ would lead to a contradiction, since the path

$$u_1v_{j-2}v_{j-3} \dots v_{j+1}v_ju_2u_3 \dots u_{m-2}u_{m-1}v_{j-1}u_m$$

would be a PCHP.

Suppose $v_k = v_{j-2}$. By (10b) we then have $c(v_{j-4}v_{j-3}) = x$. Then $n \notin \{3, 5\}$ follows from the fact that C is properly colored, and $n \notin \{4, 6\}$ holds since (11a) is not satisfied, so we deduce $n \geq 7$. Applying (4) to $s = j - 3$ and $t = j$ we have $c(v_{j-3}v_j) \neq x$. However, the path $u_1v_{j-1}u_2u_3 \dots u_{m-2}u_{m-1}v_{j-2}u_m$ and the cycle $v_jv_{j+1} \dots v_{j-4}v_{j-3}v_j$ are properly colored and satisfy (1), which contradicts our minimality assumption.

For $v_k = v_{j+2}$ we similarly conclude by (10b) and (4), with $s = j - 1$ and $t = j + 2$, that $c(v_{j-1}v_{j+2}) \neq x$. Examination of the path $u_1v_ju_2u_3 \dots u_{m-2}u_{m-1}v_{j+1}u_m$ and the

cycle $v_{j-1}v_{j+2}v_{j+3}\dots v_{j-2}v_{j-1}$ again leads to contradiction, which shows that (11a) does hold.

We have that n is an even number, and the edges of C are alternately colored x . Let $r \in \{1, 2, \dots, n\}$ be chosen so that $c(u_3v_r) \neq c(u_3u_4)$; this is possible by (3). We apply (5) and (6) with $p = 3$ and $q = r$. Then (5d) and (6d) both fail by the choice of r . Neither of the edges u_2v_{r-1} and u_2v_{r+1} have color x , due to our assumption, hence (5a) and (6a) both fail. For the same reason one of (5b) and (6b) fails, because either $v_{r-2}v_{r-1}$ or $v_{r+1}v_{r+2}$ is colored x . If $v_{r-2}v_{r-1}$ is colored x , then (5b) fails, so (5c) holds and gives $c(u_3v_r) = c(v_rv_{r+1})$, and now

$$u_1u_2v_kv_{k+1}\dots v_{r-1}v_ru_3u_4\dots u_{m-2}u_{m-1}v_{k-1}v_{k-2}\dots v_{r+2}v_{r+1}u_m$$

is a PCHP, a contradiction. If $v_{r+1}v_{r+2}$ is colored x , then (6b) fails, and $c(u_3v_r) = c(v_{r-1}v_r)$ follows similarly. But

$$u_1u_2v_{k-1}v_{k-2}\dots v_{r+1}v_ru_3u_4\dots u_{m-2}u_{m-1}v_kv_{k+1}\dots v_{r-2}v_{r-1}u_m$$

is a PCHP, again with contradiction. This finishes the proof of (12).

(13) Assume $c(u_2v_j) = x$ for some $j \in \{1, 2, \dots, n\}$, and let $w = c(v_{j-1}v_j)$ and $z = c(v_jv_{j+1})$. Then

- a)** $c(v_{j-2}v_{j-1}) = c(v_{j+1}v_{j+2}) = c(v_{j-1}v_{j+1}) = x$ and $n \in \{3, 5\}$
- b)** $c(u_2v_k) \neq x$ for all $v_k \neq v_j$
- c)** $c(u_2v_{j-1}) = c(u_2v_{j+1}) = c(u_2u_3) \in \{w, z\}$
- d)** $m \geq 4$
- e)** if $c(u_2u_3) = w$, then $c(u_3v_k) = c(u_3u_4) \neq c(u_3v_{j-1}) = w$ for all $v_k \neq v_{j-1}$.
- f)** if $c(u_2u_3) = z$, then $c(u_3v_k) = c(u_3u_4) \neq c(u_3v_{j+1}) = z$ for all $v_k \neq v_{j+1}$.

We may assume $j = 1$, so that $c(u_2v_1) = x$, $c(v_nv_1) = w$ and $c(v_1v_2) = z$. Then $c(u_2v_1) = x \neq c(u_2u_3)$ and (7b) directly imply (a). Since $n \in \{3, 5\}$ and C is properly colored, $c(u_2v_k) = x$ now implies $v_k = v_1$ by (a) and (7b), so (b) holds.

Now $c(u_2v_2) = c(u_2u_3)$ follows from (5) and (6) with $p = 2$ and $q = 2$, since (5a), (5b), (6a), (6b) all fail, and (5c) contradicts (6c), so that (5d) and the equivalent (6d) hold. Similarly we deduce $c(u_2v_n) = c(u_2u_3)$ from (5) and (6) with $p = 2$ and $q = n$.

Now (d) and $c(u_3v_1) = c(u_3u_4)$ follow from (5) and (6) with $p = 3$ and $q = 1$, as (5d) or (6d) is again satisfied. Moreover for $n = 5$ the identity $c(u_3v_3) = c(u_3u_4)$

follows, using $c(v_2v_3) \neq c(v_3v_4)$, from the fact that neither $u_1u_2v_4v_5v_1v_2v_3u_3u_4 \dots u_m$ nor $u_1v_2v_1u_2v_5v_4v_3u_3u_4 \dots u_m$ is a PCHP. Similarly we have $c(u_3v_4) = c(u_3u_4)$ when $n = 5$.

We deduce $c(u_3v_n) \in \{w, c(u_3u_4)\}$, since $u_1v_{n-1}v_{n-2} \dots v_3v_2u_2v_1v_nu_3u_4 \dots u_m$ is not a PCHP, Moreover $c(u_3v_n) \in \{c(u_2u_3), c(u_3u_4)\}$ follows, using $c(u_2v_n) = c(u_2u_3)$, by examining the path $u_1v_{n-1}v_{n-2} \dots v_2v_1u_2v_nu_3u_4 \dots u_m$, and we have $c(u_3v_n) \in \{w, c(u_3u_4)\} \cap \{c(u_2u_3), c(u_3u_4)\}$. A similar argument shows $c(u_3v_2) \in \{z, c(u_3u_4)\} \cap \{c(u_2u_3), c(u_3u_4)\}$. At least one of $c(u_3v_n)$ and $c(u_3v_2)$ is not equal to $c(u_3u_4)$, by (3), so it follows that either $c(u_3v_n) = c(u_2u_3) = w$ or $c(u_3v_2) = c(u_2u_3) = z$ holds. This shows the remaining parts of (c), (e) and (f), and (13) is proved.

(14) Assume $c(u_{m-1}v_j) = y$ for some $j \in \{1, 2, \dots, n\}$, and let $w' = c(v_{j-1}v_j)$ and $z' = c(v_jv_{j+1})$. Then

- a) $c(v_{j-2}v_{j-1}) = c(v_{j+1}v_{j+2}) = c(v_{j-1}v_{j+1}) = y$ and $n \in \{3, 5\}$
- b) $c(u_{m-1}v_k) \neq y$ for all $v_k \neq v_j$
- c) $c(u_{m-1}v_{j-1}) = c(u_{m-1}v_{j+1}) = c(u_{m-2}u_{m-1}) \in \{w', z'\}$
- d) $m \geq 4$
- e) if $c(u_{m-2}u_{m-1}) = w'$, then $c(u_{m-2}v_k) = c(u_{m-3}u_{m-2}) \neq c(u_{m-2}v_{j-1}) = w'$ for all $v_k \neq v_{j-1}$.
- f) if $c(u_{m-2}u_{m-1}) = z'$, then $c(u_{m-2}v_k) = c(u_{m-3}u_{m-2}) \neq c(u_{m-2}v_{j+1}) = z'$ for all $v_k \neq v_{j+1}$.

(14) is proved similarly to (13).

By (12) we may assume $c(u_2v_1) = x$ without loss of generality. Let $w = c(v_nv_1)$ and $z = c(v_1v_2)$. We will further assume $c(u_2v_n) = c(u_2v_2) = c(u_2u_3) = w$, which is admissible by (13c) without loss of generality. Then $m \geq 4$ holds by (13d) and

$$c(u_3v_k) = c(u_3u_4) \neq c(u_3v_n) = w \text{ for all } v_k \neq v_n$$

by (13e). These facts will be used frequently throughout the remaining part of the proof.

(15) $n = 3$.

Suppose $n \neq 3$; from (13a) it follows that n is equal to 5. Further we then have $c(v_4v_5) = c(v_2v_3) = c(v_2v_5) = x$ by (13a), and $c(u_2v_k) \neq x$ for $v_k \in \{v_3, v_4\}$ by (13b).




Figure 3: The situation in the proof of (15), where $a = c(u_3u_4)$.

We will first show that there exists $j \in \{1, 2, \dots, 5\}$ with $c(u_{m-1}v_j) = y$. So suppose not. Then by (10) there is a k such that $c(u_{m-1}v_{k-1}) = c(v_{m-1}v_k) \neq c(u_{m-2}u_{m-1})$ and $c(v_{k-2}v_{k-1}) = c(v_kv_{k+1}) = y$. The latter implies $k \notin \{1, 2\}$, using $c(v_4v_5) \neq c(v_1v_2)$ and $c(v_5v_1) \neq c(v_2v_3)$. For $k \in \{3, 4\}$ the path $u_1v_4u_2v_1v_2v_5u_3u_4 \dots u_{m-2}u_{m-1}v_3u_m$ is a PCHP, and for $k = 5$ the path $u_1v_3u_2v_1v_2v_5u_3u_4 \dots u_{m-2}u_{m-1}v_4u_m$ is a PCHP, giving a contradiction in each case. Thus we have shown that j exists as desired. Figure 3 summarizes what has been shown so far about the colors of various edges.

Now (14a) implies $y \in \{x, z, w\}$, and the value of j is uniquely determined, by (14b). Moreover, (14c) implies $c(u_{m-2}u_{m-1}) \in \{x, z, w\} \setminus \{y\}$.

Case 1: $y = x$. In this case $j = 1$, and $c(u_{m-1}v_1) = x \neq c(u_{m-2}u_{m-1})$. However, $u_1u_2v_2v_1u_{m-1}u_{m-2} \dots u_4u_3v_5v_4v_3u_m$ is a PCHP, a contradiction.

Case 2: $y = z$. In this case $j = 5$, $c(u_{m-1}v_5) = z$, and $c(u_{m-2}u_{m-1}) \in \{x, w\}$. Supposing that $c(u_{m-2}u_{m-1}) = x$ holds, the path $u_1u_2 \dots u_{m-3}u_{m-2}v_4v_3u_{m-1}u_m$ and the cycle $v_1v_5v_2v_1$ would both be properly colored, contradicting our minimality assumption. So we have $c(u_{m-2}u_{m-1}) = w$.

By (13e) we have $c(u_3v_1) = c(u_3u_4)$, and $c(u_{m-1}v_1) = c(u_{m-2}u_{m-1})$ follows from (14c), so we deduce $m \neq 4$. Since $c(u_2u_3) = w = c(u_{m-2}u_{m-1})$ it is clear that $m \neq 5$ holds, hence $m \geq 6$. By (14e) we have $c(u_{m-2}v_1) = w$. Now

$$u_1u_2v_2v_1u_{m-2}u_{m-3} \dots u_4u_3v_5v_4v_3u_{m-1}u_m$$




Figure 4: The situation in the final steps of the proof of Claim A, where $a = c(u_3u_4)$.

is a PCHP, a contradiction.

Case 3: $y = w$. In this case $j = 2$, and $c(u_{m-1}v_2) = w$. We note by (14a) that $c(v_1v_3) = w$ holds. However $u_1u_2v_4v_5u_3u_4 \dots u_m$ and $v_1v_2v_3v_1$ are now properly colored, again contradicting our minimality assumption. This finishes the last case, and (15) is proved.

We choose $j \in \{1, 2, 3\}$ with $c(u_{m-1}v_j) = y$, which is possible since (10b) fails for $n = 3$, implying that the assumption of (10) does not hold. Then j satisfies the assumption of (8), hence $c(v_{j+1}v_{j+2}) = y$ follows from (8b). We deduce $y \in \{x, z, w\}$ and proceed to divide into the three respective cases (see also Figure 4).

Case 1: $y = x$. Then $j = 1$ follows from $c(v_2v_3) = x$, and the choice of j implies $c(u_{m-1}v_1) = x$. But now $u_1u_2v_2v_1u_{m-1}u_{m-2} \dots u_4u_3v_3u_m$ is a PCHP, a contradiction.

Case 2: $y = z$. In this case $j = 3$ follows from $c(v_1v_2) = y$, implying $c(u_{m-1}v_3) = z$. By (13e) $c(u_3v_1) = c(u_3v_2) = c(u_3u_4) \neq c(u_3v_3) = w$. For $m = 4$ all of (6a,b,c,d) would fail for $p = 4$ and $q = 1$ (in particular (6b) fails by $c(u_3v_2) = c(u_3u_4) = y \neq x = c(v_2v_3)$), so we deduce $m \geq 5$.

Suppose $c(u_4v_1) \neq c(u_4u_5)$. Then $c(u_3v_1) = c(u_4v_1)$ holds, as otherwise the path $u_1u_2v_2v_3u_3v_1u_4u_5 \dots u_m$ would be a PCHP. Now (6) with $p = 4$ and $q = 1$ implies $c(u_3v_2) = x$ or $c(u_4v_1) = w$ (i.e. (6b) or (6c)). We have $c(u_4v_1) = c(u_3v_1) = c(u_3u_4) \neq$

$c(u_2u_3) = w$, so $c(u_3v_2) = x$ follows. But then $u_1u_2v_2u_3v_3u_{m-1}u_{m-2}\dots u_5u_4v_1u_m$ is a PCHP, which is a contradiction. We conclude that $c(u_4v_1) = c(u_4u_5)$ holds.

Suppose $c(u_4v_2) \neq c(u_4u_5)$. Then (5c) for $p = 4$ and $q = 2$ holds, that is $c(u_4v_2) = x$, and $u_1u_2u_3v_1v_3u_{m-1}u_{m-2}\dots u_5u_4v_2u_m$ is a PCHP, a contradiction. Hence $c(u_4v_2) = c(u_4u_5)$ follows. Now by (3) we have $c(u_4v_3) \neq c(u_4u_5)$. With $p = 4$ and $q = 3$ either (5b) or (5c) holds, hence $c(u_3v_2) = z$ or $c(u_4v_3) = w$. Examination of the path $u_1v_2u_2v_1u_3v_3u_4u_5\dots u_m$ allows us to deduce $c(u_3v_1) = x$ or $c(u_4v_3) = w$. Then $c(u_4v_3) = w$ follows, since the two alternatives conflict due to $c(u_3v_1) = c(u_3v_2)$.

Let i be the largest number such that $i \leq m$ and such that $c(u_{i'}v_1) = c(u_{i'}v_2) = c(u_{i'}u_{i'+1}) \neq c(u_{i'}v_3) = w$ holds for every $i' = 3, 4, \dots, i - 1$. We note that i exists and satisfies $i \geq 5$. It is useful to observe that (5a) and (6a) fail for $p = i$ and every $q = 1, 2, 3$; for (5a) this follows from $c(u_{i-2}u_{i-1}) \neq w$ for $q = 1$, and it follows from $c(u_{i-1}v_{q-1}) = c(u_{i-1}u_i) \neq c(u_{i-2}u_{i-1})$ for $q \neq 1$. Similarly for (6a).

For $i = m$ all of (6a,b,c,d) fail for $p = i$ and $q = 1$ (in particular (6b) fails by $c(u_{i-1}v_2) = c(u_{i-1}u_i) = y \neq x = c(v_2v_3)$), so we deduce $i < m$.

Suppose $c(u_i v_1) \neq c(u_i u_{i+1})$. For $p = i$ and $q = 1$ (5) and (6) imply $c(u_i v_1) = z$ and $c(u_{i-1} v_2) = x$ (respectively (5c) and (6b)). Using $c(u_{i-1} v_1) = c(u_{i-1} v_2) = x$ and $c(u_{i-2} v_2) = c(u_{i-2} u_{i-1}) \neq c(u_{i-1} u_i) = c(u_{i-1} v_2) = x$, the path

$$u_1 \dots u_{i-3} u_{i-2} v_2 v_3 u_{i-1} v_1 u_i u_{i+1} \dots u_m$$

is a PCHP. This contradiction shows $c(u_i v_1) = c(u_i u_{i+1})$. Suppose $c(u_i v_2) \neq c(u_i u_{i+1})$. Then (5) with $p = i$ and $q = 2$ implies $c(u_i v_2) = x$, but

$$u_1 u_2 \dots u_{i-2} u_{i-1} v_1 v_3 u_{m-1} u_{m-2} \dots u_{i+1} u_i v_2 u_m$$

is a PCHP, a contradiction. Therefore $c(u_i v_2) = c(u_i u_{i+1})$. By (3) we have $c(u_i v_3) \neq c(u_i u_{i+1})$. Then $c(u_i v_3) \neq w$ follows from the choice of i , and

$$u_1 v_2 v_1 u_2 u_3 \dots u_{i-2} u_{i-1} v_3 u_i u_{i+1} \dots u_m$$

is a PCHP, a contradiction.

Case 3: $y = w$. This implies $j = 2$ and $c(u_{m-1} v_2) = w$. By (14c) we have $c(u_{m-2} u_{m-1}) \in \{x, z\}$. The case $c(u_{m-2} u_{m-1}) = z$ is symmetric to Case 2, so only the case $c(u_{m-2} u_{m-1}) = x$ remains. Then $m \geq 5$ follows, as P is properly colored. We can assume $c(u_2 u_{m-1}) \neq x$ without loss of generality. But $u_1 u_2 u_{m-1} u_{m-2} \dots u_4 u_3 v_3 v_2 v_1 u_m$ is a PCHP, with contradiction. This finishes the proof of Case 3, and of Claim A. \square

References

- [1] N. Alon and G. Gutin, Properly colored Hamilton cycles in edge colored complete graphs. *Random Structures and Algorithms* **11** (1997) 179-186.
- [2] J. Bang-Jensen and G. Gutin, Alternating cycles and paths in edge-coloured multi-graphs: a survey. *Discrete Math.* **165/166** (1997) 39–60.
- [3] J. Bang-Jensen and G. Gutin, *Digraphs: Theory, Algorithms and Applications*, Springer-Verlag, London, 2000.
- [4] J. Bang-Jensen, G. Gutin and A. Yeo, A polynomial algorithm for the Hamiltonian cycle problem in semicomplete multipartite digraphs. *J. Graph Theory* **29** (1998) 111-132.
- [5] J. Bang-Jensen, G. Gutin and A. Yeo, Properly coloured Hamiltonian paths in edge-coloured complete graphs. *Discrete Appl. Math.* **82** (1998) 247-250.
- [6] M. Bankfalvi and Zs. Bankfalvi, Alternating hamiltonian circuit in two-coloured complete graphs, (Proc. Colloq. Tihany 1968), Academic Press, New York (1968) 11-18.
- [7] O. Barr, Properly coloured Hamiltonian paths in edge-coloured complete graphs without monochromatic triangles. *Ars Combinatoria* **50** (1998) 316-318.
- [8] A. Benkouar, Y. Manoussakis, V. Paschos and R. Saad, On the complexity of finding alternating Hamiltonian and Eulerian cycles in edge-coloured graphs. Lecture Notes in Comput. Sci., **557** (Springer, Berlin, 1991) 190-198.
- [9] B. Bollobás and P. Erdős, Alternating Hamiltonian cycles. *Israel J. Math.* **23** (1976) 126-131.
- [10] G. Gutin, Finding a longest path in a complete multipartite digraph. *SIAM J. Discrete Math.* **6** (1993) 270-273
- [11] R. Saad, Finding a longest alternating cycle in a 2-edge-coloured complete graph is in RP. *Combinatorics, Probability and Computing* **5** (1996) 297-306.
- [12] J. Shearer, A property of the colored complete graph. *Discrete Math.* **25** (1979) 175-178.